

## **Massive Galaxies at High Redshift from K20+GOODS**

**Alvio Renzini**, European Southern Observatory, ESO

**Abstract** With the aim of tracing the assembly of massive galaxies through cosmic time, spectroscopic redshifts have been measured for a sample of K-band luminous galaxies ( $K_s < 20$ ) at  $1.7 < z < 2.3$ , selected in a field covered by both the K20 and the GOODS Surveys. Fitting of their multi-color spectral energy distributions indicates stellar masses  $M \sim 10^{11}$  Msun or more for most of the program galaxies, while extinction corrected star formation rates (SFRs) of 100--500 Msun/yr are derived. Their rest-frame UV morphologies from HST/GOODS imaging are highly irregular, suggesting that merging-driven starbursts are going on in these galaxies. Morphologies tend to be more compact in the near-IR, a hint for the possible presence of older stellar populations. Such galaxies are strongly clustered, with the majority belonging to redshift spikes, which indicates a correlation length  $r_o = (9-17) h$  Mpc (1 sigma range). Current semianalytical models of galaxy formation appear to underpredict by more than an order of magnitude the number density of such a population of massive and powerful starburst galaxies at  $z=2$ . The high masses and SFRs together with the strong clustering suggest that at  $z=2$  we may have started to explore the major formation epoch of massive early-type galaxies.